

Supplemental material

Presence of *Malassezia* hyphae is correlated with pathogenesis of seborrheic dermatitis

Juanjuan Li^{1#}, Yahui Feng^{1#}, Chen Liu², Zhiya Yang², Sybren de Hoog³, Yuying Qu¹, Biao Chen², Dongmei Li⁴, Huabao Xiong^{5*}, Dongmei Shi^{2,6*}

¹Department of Clinical Medicine, Jining Medical University, Jining 272067, Shandong, China

²Laboratory of Medical Mycology, Jining No. 1 People's Hospital, Jining 272067, Shandong, China

³Centre of Expertise in Mycology of Radboud University Medical Centre / Canisius Wilhelmina Hospital, 6525, Nijmegen, The Netherlands

⁴Department of Microbiology & Immunology, Georgetown University Medical Center, Washington DC 20057, USA

⁵Institute of Immunology and Molecular Medicine, Basic Medical School, Jining Medical University, Jining 272067, China.


⁶Department of Dermatology, Jining No.1 People's Hospital, Jining 272067, Shandong, China

The authors contributed to this article equally.

Corresponding authors:

Correspondence to: Dongmei Shi, Dermatology Department of the First People's Hospital of Jining, Shandong, 272011, China. Phone: +86 537-6051008; Email: shidongmei28@163.com, or to Huabao Xiong, Institute of Immunology and Molecular Medicine, Jining Medical University, Jining, Shandong, 272067, China, Phone: +86 537-3616283; Email: xionghbl@yahoo.com

Supplemental Fig. 1



Supplemental Figure 1. A: General information of Hyphae-positive groups and Hyphae-negative groups. B: Age distribution of Hyphae-positive group and Hyphae-negative group.